

ON TRANSLATION INVARIANT SYMMETRIC POLYNOMIALS AND HALDANE'S CONJECTURE

JESSE LIPTRAP

ABSTRACT. We show that the ring of translation invariant symmetric polynomials in n variables is isomorphic to the full polynomial ring in $n - 1$ variables, in characteristic 0. We disprove a conjecture of Haldane regarding the structure of such polynomials. Our motivation is the fractional quantum Hall effect, where translation invariant (anti)symmetric complex n -variate polynomials characterize n -electron wavefunctions.

1. INTRODUCTION

A polynomial $p(z_1, \dots, z_n)$ is *translation invariant* if

$$p(z_1 + c, \dots, z_n + c) = p(z_1, \dots, z_n)$$

for all c . Over $\mathbb{C}$, such a polynomial might yield a quantum mechanical description of n particles in the plane. Such is the nature of fractional quantum Hall states, where in addition our polynomials are symmetric or antisymmetric (Wen and Wang [5]). Thus we are led to the study of translation invariant (anti)symmetric polynomials. Antisymmetric polynomials fortunately do not need special treatment, as they are merely symmetric polynomials multiplied by the Vandermonde determinant $\prod_{i < j} (z_i - z_j)$.

This elementary note contains two main results on the structure of translation invariant symmetric polynomials. First, a simple description of the ring of all such polynomials (Corollary 2.3). Second, a counterexample to Haldane's conjecture [3] that every homogeneous translation invariant symmetric polynomial satisfies a certain physically convenient property (Proposition 3.7). More precisely, each symmetric polynomial p is associated with a finite poset $B(p)$; Haldane conjectured that if p is homogeneous and translation invariant, then $B(p)$ has a maximum. We prove the conjecture for polynomials of at most three variables, construct a minimal counterexample, and discuss whether a weakened version of the conjecture holds.

2. THE RING OF TRANSLATION INVARIANT SYMMETRIC POLYNOMIALS

Throughout, let $\mathbb{F}$ be a field of characteristic 0. We begin with just translation invariance. Let $T \subseteq \mathbb{F}[z_1, \dots, z_n]$ be the ring of translation invariant polynomials. Imagining $z_1, \dots, z_n$ as the locations of n identical particles, and $x_1, \dots, x_n$ as the corresponding center of mass coordinates, our main theorem says that T written in terms of $x_1, \dots, x_n$ is $\mathbb{F}[x_1, \dots, x_n]$ modulo one degree of freedom.

Theorem 2.1. *Let $\rho: \mathbb{F}[x_1, \dots, x_n] \rightarrow T$ be the surjective algebra homomorphism*

$$x_i \mapsto z_i - z_{\text{avg}}$$

where $z_{\text{avg}} = \frac{1}{n}(z_1 + \dots + z_n)$. Then $\ker \rho = (x_{\text{avg}})$.

Proof. Section 4. □

Now let $R \subseteq T$ be the ring of translation invariant symmetric polynomials in $z_1, \dots, z_n$, and $S \subseteq \mathbb{F}[x_1, \dots, x_n]$ be the ring of symmetric polynomials in $x_1, \dots, x_n$.

Corollary 2.2. *Let $\sigma: S \rightarrow R$ agree with ρ . Then σ is a surjective algebra homomorphism, with kernel $(x_1 + \dots + x_n)$.*

Proof. It suffices to show $\rho(S) = R$. Clearly $\rho(S) \subseteq R$. Given $p(z_1, \dots, z_n) \in R$, translation invariance yields

$$p(z_1, \dots, z_n) = p(z_1 - z_{\text{avg}}, \dots, z_n - z_{\text{avg}}) = \rho(p(x_1, \dots, x_n)).$$

Thus $\rho(S) = R$. □

Since $\mathbb{F}$ has characteristic 0, any element of S can be written uniquely as a polynomial in the power sum symmetric polynomials $x_1^k + \dots + x_n^k$, where $1 \leq k \leq n$. In other words, the algebra homomorphism $\theta: \mathbb{F}[w_1, \dots, w_n] \rightarrow S$ defined by $\theta(w_k) = x_1^k + \dots + x_n^k$ is an isomorphism. Note that we could define a different isomorphism θ using elementary symmetric polynomials or complete homogeneous symmetric polynomials. In any case, $\sigma\theta: \mathbb{F}[w_1, \dots, w_n] \rightarrow R$ is a surjective algebra homomorphism, with kernel (w_1) .

Corollary 2.3. *The algebra homomorphism $\mathbb{F}[w_2, \dots, w_n] \rightarrow R$ given by*

$$w_k \mapsto (z_1 - z_{\text{avg}})^k + \dots + (z_n - z_{\text{avg}})^k$$

is an isomorphism.

Next we consider the vector space R^d of all polynomials in R which are homogeneous of degree d . Let f be the above isomorphism. Since $f(w_k)$ is homogeneous of degree k , we obtain a basis for R^d , namely all

$$(1) \quad w_\lambda = \prod_{k=2}^n f(w_k)^{\lambda_k}$$

where λ is any partition of d into integers between 2 and n , and λ_k is the multiplicity of k in λ . Simon, Rezayi, and Cooper [4] prove directly that these w_λ form a basis of R^d , whereas we have deduced this fact from the ring structure of R . Although [4] defines w_λ using elementary symmetric polynomials rather than power sum symmetric polynomials, this difference is purely cosmetic. Since the dimension m_d of R^d is the number of partitions of d into integers between 2 and n , a generating function for m_d is

$$\sum_{d=0}^{\infty} m_d t^d = \prod_{s=2}^n \frac{1}{1 - t^s}$$

Finally, we describe the vector space $A \subset \mathbb{F}[z_1, \dots, z_n]$ of translation invariant antisymmetric polynomials. It is well-known that any antisymmetric polynomial can be written uniquely as $q\Delta$, where q is a symmetric polynomial and Δ is the Vandermonde determinant $\prod_{i < j} (z_i - z_j)$. Since Δ is translation invariant, we have $A = R\Delta$, defining a vector space isomorphism $R \rightarrow A$, which sends each basis (1) to a basis for the vector space of homogeneous translation invariant antisymmetric polynomials of degree $d + n(n-1)/2$.

3. HALDANE'S CONJECTURE

Every symmetric polynomial is a unique linear combination of symmetrized monomials, which physicists like to call *boson occupation states*. We identify symmetrized monomials with multisets of natural numbers:

$$[l_1, \dots, l_n] = \sum_{\sigma \in \text{Sym}(n)} z_{\sigma(1)}^{l_1} \cdots z_{\sigma(n)}^{l_n}$$

For instance, the multiset $[5, 0, 0]$ corresponds to the symmetrized monomial $2z_1^5 + 2z_2^5 + 2z_3^5$. *Squeezing* a symmetrized monomial $[l_1, \dots, l_n]$ means decrementing l_i and incrementing l_j for any pair of indices i, j such that $l_i > l_j + 1$. The *squeezing order* is a partial order on symmetrized monomials: put $s > t$ iff t can be obtained from s by repeated squeezing. For a symmetric polynomial p , let $B(p)$ be the set of all symmetrized monomials with nonzero coefficient in p . We view $B(p)$ as a poset under the squeezing order and refer to it as the *squeezing poset* of p .

Definition 3.1. A symmetric polynomial is *Haldane* if its squeezing poset has a maximum.

Conjecture 3.2 (Haldane [3]). *Every homogeneous translation invariant symmetric polynomial is Haldane.*

Remark. Since squeezing preserves homogeneous degree, Haldane polynomials are homogeneous. Many homogeneous symmetric polynomials are not Haldane, such as $[3, 3, 0] + [4, 1, 1]$, but these might not be translation invariant.

Proposition 3.3. *Haldane's conjecture holds for polynomials of ≤ 3 variables.*

Proof. The conjecture is vacuously true for univariate polynomials. Every bivariate symmetrized monomial of homogeneous degree d has the form $[a, b]$, with $a + b = d$. These are linearly ordered under squeezing, so Haldane's conjecture is automatic in the bivariate case.

For the trivariate case, define $\tau: \mathbb{F}[z_1, z_2, z_3] \rightarrow \mathbb{F}[z_1, z_2, z_3, t]$ by

$$\tau(p)(z_1, z_2, z_3, t) = p(z_1 + t, z_2 + t, z_3 + t),$$

so that p is translation invariant iff $\tau(p) = p$. Define linear endomorphisms τ_i of $\mathbb{F}[z_1, z_2, z_3]$ by $\tau(p) = \sum_{i=0}^d \tau_i(p)t^i$, so that p is translation invariant iff $\tau_i(p) = 0$ for all $i > 0$. Then

$$\tau_1([a, b, c]) = a[a - 1, b, c] + b[a, b - 1, c] + c[a, b, c - 1]$$

for all $a, b, c > 0$. Now suppose $[a, b, c]$ is a maximal element of the squeezing poset of some $p \in R_3^d$, with $a \geq b \geq c > 0$. Then $[a + 1, b, c - 1]$ and $[a, b + 1, c - 1]$ are not in $B(p)$. The above equation then implies that the coefficient of $[a, b, c]$ in p equals c times the coefficient of $[a, b, c - 1]$ in $\tau_1(p)$. Thus $\tau_1(p) \neq 0$, contradicting the translation invariance of p . Therefore every maximal element of $B(p)$ has the form $[a, b, 0]$, with $a + b = d$. These are linearly ordered under squeezing; their maximum is the maximum of $B(p)$. $\square$

Any two symmetrized monomials written as weakly decreasing sequences of natural numbers can be compared lexicographically. The lexicographic order $>_{\text{lex}}$ on symmetrized monomials linearizes the squeezing order. Let R_n^d be the vector space of translation invariant symmetric n -variate polynomials of homogeneous degree d ,

and let L_n^d be the set of lexicographic maxima of squeezing posets of polynomials in R_n^d . Note $|L_n^d| \leq \dim R_n^d$.

Definition 3.4. A symmetrized monomial s is *completely squeezable* if $s >_{\text{lex}} t$ implies $s > t$, for all symmetrized monomials t .

Lemma 3.5. *If every element of L_n^d is completely squeezable, then Haldane's conjecture holds for R_n^d . If Haldane's conjecture holds for R_n^d , then L_n^d is linearly ordered under squeezing.*

Proof. The first statement is immediate. For the second, suppose $m_1, m_2 \in L_n^d$ are incomparable. Let $p_1, p_2 \in R_n^d$ such that m_i is the lexicographic maximum of $B(p_i)$ for $i = 1, 2$. W.l.o.g. assume m_1 is lexicographically bigger than m_2 , and let c_i be the coefficient of m_2 in p_i . Choose a scalar $c \neq -c_1/c_2$, and let $q = p_1 + cp_2$. Then $q \in R_n^d$ and $m_1, m_2 \in B(q)$. Since m_1 is the lexicographic maximum of $B(q)$, it is maximal in $B(q)$ under squeezing. Since m_1 and m_2 are incomparable, q is not Haldane. $\square$

Proposition 3.6. *Haldane's conjecture holds for R_4^d with $d < 14$ but fails for R_4^{14} .*

Proof. It is a straightforward computational linear algebraic exercise to compute L_n^d using the basis for R_n^d given by formula (1). Since every symmetrized monomial

d	L_4^d
0	$\emptyset$
1	$\emptyset$
2	$\{[2, 0, 0, 0]\}$
3	$\{[3, 0, 0, 0]\}$
4	$\{[4, 0, 0, 0], [2, 2, 0, 0]\}$
5	$\{[5, 0, 0, 0]\}$
6	$\{[6, 0, 0, 0], [4, 2, 0, 0], [3, 3, 0, 0]\}$
7	$\{[7, 0, 0, 0], [5, 2, 0, 0]\}$
8	$\{[8, 0, 0, 0], [6, 2, 0, 0], [5, 3, 0, 0], [4, 4, 0, 0]\}$
9	$\{[9, 0, 0, 0], [7, 2, 0, 0], [6, 3, 0, 0]\}$
10	$\{[10, 0, 0, 0], [8, 2, 0, 0], [7, 3, 0, 0], [6, 4, 0, 0], [5, 5, 0, 0]\}$
11	$\{[11, 0, 0, 0], [9, 2, 0, 0], [8, 3, 0, 0], [7, 4, 0, 0]\}$
12	$\{[12, 0, 0, 0], [10, 2, 0, 0], [9, 3, 0, 0], [8, 4, 0, 0], [7, 5, 0, 0], [6, 6, 0, 0], [6, 4, 2, 0]\}$
13	$\{[13, 0, 0, 0], [11, 2, 0, 0], [10, 3, 0, 0], [9, 4, 0, 0], [8, 5, 0, 0]\}$
14	$\{[14, 0, 0, 0], [12, 2, 0, 0], [11, 3, 0, 0], [10, 4, 0, 0], [9, 5, 0, 0], [8, 6, 0, 0], [8, 4, 2, 0], [7, 7, 0, 0]\}$

TABLE 1. Enumeration of lexicographic maxima.

of the form $[a, b, 0, \dots, 0]$ is completely squeezable, as is $[6, 4, 2, 0]$, we see that every element of L_4^d , $d < 14$, is completely squeezable (Table 1). But L_4^{14} is not linearly ordered under squeezing: $[8, 4, 2, 0]$ and $[7, 7, 0, 0]$ are incomparable. Then apply Lemma 3.5. $\square$

It is a straightforward computational linear algebraic exercise to construct a non-Haldane polynomial in R_4^{14} by following the proof of Lemma 3.5. We get

$$\begin{aligned} p = & 3[8, 4, 2, 0] - 3[8, 4, 1, 1] - 3[8, 3, 3, 0] + 6[8, 3, 2, 1] - 3[8, 2, 2, 2] \\ & + 3[7, 7, 0, 0] - 42[7, 6, 1, 0] + 46[7, 5, 2, 0] + 80[7, 5, 1, 1] - 22[7, 4, 3, 0] \\ & - 188[7, 4, 2, 1] + 112[7, 3, 3, 1] + 8[7, 3, 2, 2] + 77[6, 6, 2, 0] + 70[6, 6, 1, 1] \\ & - 182[6, 5, 3, 0] - 700[6, 5, 2, 1] + 112[6, 4, 4, 0] + 168[6, 4, 3, 1] + 1078[6, 4, 2, 2] \\ & - 728[6, 3, 3, 2] + 5[5, 5, 4, 0] + 1072[5, 5, 3, 1] + 246[5, 5, 2, 2] - 722[5, 4, 4, 1] \\ & - 2976[5, 4, 3, 2] + 1808[5, 3, 3, 3] + 1805[4, 4, 4, 2] - 1130[4, 4, 3, 3]. \end{aligned}$$

Proposition 3.7. *The polynomial p is a minimal counterexample to Haldane's conjecture.*




FIGURE 1. Hasse diagram of $B(p)$. Arrows point from smaller to bigger elements.

Proof. One checks by computer that p is translation invariant. Since it is symmetric and homogeneous but lacks a maximum (Figure 1), it breaks Haldane's conjecture. By Proposition 3.6 it is minimal with respect to arity and homogeneous degree. $\square$

Remark. The counterexample of Proposition 3.7 is not minimal with respect to homogeneous degree. For instance, the smallest pentavariate counterexamples have homogeneous degree 10.

Remark. We might weaken Haldane's conjecture by hoping R_n^d has a basis of Haldane polynomials. Computer evidence suggests $|L_n^d| = \dim R_n^d$. Writing $L_n^d = \{l_1, \dots, l_k\}$, we could then obtain a special basis $\{p_1, \dots, p_k\}$ of R_n^d satisfying

$B(p_i) \cap L_n^d = \{l_i\}$. Perhaps it would be a Haldane basis or could be used to construct one.

4. PROOF OF THEOREM 2.1

We factor ρ into two maps which are easier to study:

$$\begin{array}{ccccc} & & \rho & & \\ & \nearrow & & \searrow & \\ X & \xrightarrow{\tau} & Y & \xrightarrow{\pi} & T \end{array}$$

Let $Y = \mathbb{F}[y_1, \dots, y_n]$, and define τ, π by

$$\tau(x_i) = \frac{1}{n} \sum_{j=0}^{n-1} (n-1-j)y_{i+j} \quad \pi(y_i) = z_i - z_{i+1}$$

where index addition is modulo n . Then

$$\pi\tau(x_i) = \frac{1}{n} \left((n-1)z_i - \sum_{j \neq i} z_j \right) = z_i - z_{\text{avg}}$$

showing $\pi\tau = \rho$. It suffices to show $\tau^{-1}(\ker \pi) = (x_{\text{avg}})$. Since $\tau(x_{\text{avg}}) \propto y_{\text{avg}}$, this follows from Lemmas 4.1 and 4.3 below.

Lemma 4.1. τ is an isomorphism.

Proof. Let $\hat{\tau}: \mathbb{F}x_1 + \dots + \mathbb{F}x_n \rightarrow \mathbb{F}y_1 + \dots + \mathbb{F}y_n$ be the linear map which extends to τ . Then the matrix M of $\hat{\tau}$ with respect to the evident bases is the $n \times n$ circulant matrix with first column vector

$$v = \frac{1}{n}(n-1, n-2, \dots, 0).$$

Then $M^\top$ is the circulant matrix with first row v . Since $\text{char } \mathbb{F} = 0$, the entries of v form a strictly decreasing sequence of nonnegative reals. Therefore $M^\top$ is nonsingular by Theorem 3 of Geller, Kra, Popescu, and Simanca [2]. Hence $\hat{\tau}$ is an isomorphism. Therefore τ is an isomorphism by Observation 4.2. $\square$

Observation 4.2. Suppose $f: \mathbb{F}[a_1, \dots, a_n] \rightarrow \mathbb{F}[b_1, \dots, b_n]$ is an algebra homomorphism between polynomial rings which restricts to a linear map

$$\hat{f}: \mathbb{F}a_1 + \dots + \mathbb{F}a_n \rightarrow \mathbb{F}b_1 + \dots + \mathbb{F}b_n$$

If $\hat{f}$ is an isomorphism, then so is f .

Proof. The universal property of polynomial rings. $\square$

Lemma 4.3. $\ker \pi = (y_{\text{avg}})$.

Proof. Let $\alpha = (\alpha_1, \alpha_2, \alpha_3)$ be the chain map

$$\begin{array}{ccccccc} 0 & \longrightarrow & (y_{\text{avg}}) & \longrightarrow & Y & \xrightarrow{\pi} & T \longrightarrow 0 \\ & & \alpha_1 \uparrow & & \alpha_2 \uparrow & & \alpha_3 \uparrow \\ 0 & \longrightarrow & (y_1) & \longrightarrow & Y & \xrightarrow{\pi'} & Y' \longrightarrow 0 \end{array}$$

where $Y' = \mathbb{F}[y_2, \dots, y_n]$, and $\pi', \alpha_3, \alpha_2, \alpha_1$ are the algebra homomorphisms such that π' kills y_1 and fixes the other variables, α_3 and π agree, α_2 sends y_1 to y_{avg}

and fixes the other variables, α_1 and α_2 agree, and the unlabelled nonzero maps are inclusions of ideals. We want the top sequence to be exact. Since the bottom sequence is exact, it suffices to check α is a chain isomorphism.

Since α is a chain map, it suffices to show each component is an isomorphism. By Observation 4.2, α_2 is an isomorphism. Then so is α_1 . For α_3 , let $\beta: Y' \rightarrow Y'$ be the algebra homomorphism given by $\beta(y_i) = y_i + \cdots + y_n$ for $2 \leq i \leq n$. Again by Observation 4.2, β is an isomorphism. Then it suffices to show $\gamma = \alpha_3\beta$ is an isomorphism.

Note $\gamma: Y' \rightarrow T$ and $\gamma(y_i) = z_i - z_1$ for $2 \leq i \leq n$. Since

$$p(z_1, \dots, z_n) = p(0, z_2 - z_1, \dots, z_n - z_1) = \gamma(p(0, y_2, \dots, y_n))$$

for any $p(z_1, \dots, z_n) \in T$, the homomorphism γ is surjective. If

$$0 = \gamma(q(y_2, \dots, y_n)) = q(z_2 - z_1, \dots, z_n - z_1)$$

then $0 = \pi'(q(y_2 - y_1, \dots, y_n - y_1)) = q(y_2, \dots, y_n)$, showing γ is injective. Thus γ is an isomorphism. $\square$

ACKNOWLEDGEMENTS

It is a pleasure to thank Tobias Hagge, Thomas Howard, Joules Nahas, and Zhenghan Wang for useful discussions, and Zhenghan Wang for suggesting the two problems of this paper.

REFERENCES

- [1] David Amos. *Haskell for Maths (Version 1)*, 2008. <http://www.polyomino.f2s.com/david/haskell/main.html>.
- [2] Daryl Geller, Irwin Kra, Sorin Popescu, and Santiago Simanca. On circulant matrices. www.math.sunysb.edu/~sorin/eprints/circulant.pdf.
- [3] F. Haldane. *Bull. Am. Phys. Soc.*, 51:633, 2006. Conference talk.
- [4] Steven H. Simon, E. H. Rezayi, and Nigel R. Cooper. Pseudopotentials for multi-particle interactions in the quantum Hall regime. *Phys. Rev. B*, 75(195306), 2007, cond-mat/0701260.
- [5] Xiao-Gang Wen and Zhenghan Wang. Classification of symmetric polynomials of infinite variables: Construction of abelian and non-abelian quantum Hall states. *Phys. Rev. B*, 77(23):235108, Jun 2008, arXiv:0801.3291v2.

MATHEMATICS DEPARTMENT, UNIVERSITY OF CALIFORNIA, SANTA BARBARA, CA 93106, U.S.A.
E-mail address: jliptrap@math.ucsb.edu